



Short communication

Phytomyxid infection in the non-native seagrass *Halophila stipulacea* in St Eustatius, Caribbean Netherlands

Anna Maitz^{a,b,*}, Kimani Kitson-Walters^{a,b}, Erik Maitz Boman^c

^a Caribbean Netherlands Science Institute (CNSI), P.O. Box 65, St. Eustatius, Caribbean Netherlands

^b NIOZ Royal Netherlands Institute for Sea Research, Utrecht University, P.O. Box 59, 1790 AB Den Burg, Texel, the Netherlands

^c St Eustatius National Parks Foundation (STENAPA), Oranjebaai, St Eustatius, Caribbean Netherlands

ARTICLE INFO

Keywords:

Non-native seagrass
Plant parasite
Aquatic plant
Infection
Gall
Morphological change

ABSTRACT

Phytomyxids are a monophyletic group of biotrophs/parasites of a variety of organisms including seagrasses with a wide distribution range that includes the Caribbean. The seagrass *Halophila stipulacea*, native to the Indo-Pacific and Red Sea, is a known host for phytomyxids in the Mediterranean. However, to date phytomyxid infection has not been reported for *H. stipulacea* in the Caribbean. Infection in *H. stipulacea* is characterized by swelling of the leaf petioles due to gall formation, and coloration of these galls varies depending on the stage of maturity. *H. stipulacea* fragments with an apparent phytomyxid infection as well as uninfected fragments were collected in St Eustatius, north-eastern Caribbean, for comparative biometric analysis. Measurements of leaf length, leaf width, internode and root length were taken. Infected *H. stipulacea* fragments were significantly smaller than uninfected fragments across all biometrics measured, and exhibited similar gall colorations and swelling of the leaf petioles previously described for *H. stipulacea* in the Mediterranean. Based on our observations, the apparent infection in *H. stipulacea* fragments on St. Eustatius is likely caused by a phytomyxid parasite and is the first record of phytomyxid infection of this seagrass species in the Caribbean.

1. Introduction

Phytomyxids are a monophyletic group of obligate intracellular biotrophs/parasites which utilize a variety of host species (e.g. green plants, diatoms, brown algae, stramenopiles) and are found in many freshwater, marine and terrestrial ecosystems (Braselton, 1995; Maier et al., 2000; Neuhauser et al., 2011). In marine ecosystems, seagrasses are known to be susceptible to phytomyxid infections (Den Hartog, 1989; Karling, 1968; Braselton and Short, 1985). Depending on the species of seagrass and pathogen, seagrasses infected by phytomyxids may form characteristic galls in the leaf petioles (Den Hartog, 1989; Vohnik et al., 2017).

Phytomyxids are present in both temperate (4–24 °C) and tropical (>24 °C) regions (Den Hartog, 1989), including the wider Caribbean Region (Walker and Campbell, 2009). Several seagrass genera are compatible host species for phytomyxids e.g. *Zostera* spp., *Ruppia* spp., *Halodule* spp., *Halophila* spp. (Den Hartog, 1989; Ferdinandsen and Winge, 1914; Karling, 1968; Vohnik et al., 2017). *H. stipulacea* is native to the Indian Ocean and Red Sea (Den Hartog, 1970) and has spread rapidly throughout the eastern and southern Caribbean following its

initial observation in 2002 (Ruiz and Ballantine, 2004). Phytomyxid infection (*Plasmodiophora diplantherae*) has previously been found in a native Caribbean seagrass species (*Halodule wrightii*; Ferdinandsen and Winge, 1914; Walker and Campbell, 2009). However, to our knowledge, phytomyxid infection in *H. stipulacea* has to date not been recorded in the Caribbean. Here, we report the first observation of apparent phytomyxid infection in *H. stipulacea* in St. Eustatius. Our results increase knowledge about phytomyxid distribution, as well as the morphological effects on infected seagrass.

2. Methods

H. stipulacea fragments with apparent phytomyxid infection were discovered in St Eustatius, Caribbean Netherlands, at location 17.479867° -62.994017° in October 2018 at a depth of 18 m. During three subsequent exploration dives using SCUBA at the same location on 28th February 2020 and 2nd and 3rd March 2020, two divers identified and collected nine infected *H. stipulacea* fragments. In addition, *H. stipulacea* fragments (n = 9) with no apparent sign of infection were randomly collected immediately after and within 1 m of the infected

* Corresponding author at: Caribbean Netherlands Science Institute (CNSI), P.O. Box 65, St. Eustatius, Caribbean Netherlands.

E-mail address: anna.maitz@nioz.nl (A. Maitz).

<https://doi.org/10.1016/j.aquabot.2020.103321>

Received 1 September 2020; Received in revised form 6 October 2020; Accepted 9 October 2020

Available online 14 October 2020

0304-3770/© 2020 Elsevier B.V. All rights reserved.




Fig. 1. Left image (A): Infected *H. stipulacea* with black galls in the petiole of the shoot. Leaves have stunted growth; rhizomes and roots are also shorter than uninfected seagrass. Right image (B): Uninfected *H. stipulacea* fragment as comparison found on St Eustatius, Caribbean Netherlands.




Fig. 2. Left image (A) shows the infection in different stages by color difference from white, light brown, ochre and black galls in the leaf petiole. Right image (B) shows an enlarged version of color variation and structure of the galls in *H. stipulacea* found on St Eustatius, Caribbean Netherlands.

fragment and used as comparative controls. The healthy fragments were carefully collected to contain all morphological parts of the seagrass (i.e. leaves, internodes and roots). The infected fragments collected were limited to the section of the seagrass that appeared to be affected. All fragments were collected by hand and placed in individual sample bags and subsequently brought to a laboratory for biometric analysis. Measurements (mm) of leaf length, leaf width, internode length, and root length were recorded for each fragment. Identification of a phytomyxid infection in *H. stipulacea* on St. Eustatius was based on a literature review and images by Vohnik et al. (2017).

Statistical analyses were conducted in the R software environment version 4.0.0 (R Core Team, 2020), using the packages *car* (Fox and Weisberg, 2018), and *dplyr* (Wickham, 2018). Measurements from all fragments were combined for biometric comparison between infected and uninfected fragments ($n = 9$). Normality and homogeneity of variance between the groups were tested with a Shapiro-Wilk (Royston, 1995) and Levene's test (Levene, 1960). When normality and homogeneity of variance could be assumed, which was always the case, a Student's *t*-test was performed to test for differences between the biometrics of the infected and uninfected fragments.

3. Results and discussion

Gall formation was visible in the leaf petioles of the infected

H. stipulacea fragments, where gall coloration ranged from black, brown, and/or beige to off-white. According to Kolátková et al. (2020) and Vohnik et al. (2017), the color differences represent the developmental stages of the infection. White colored galls contain sporogenic plasmodia representing early infection (Fig. 1A), and dark colored galls indicate mature resting spores (Fig. 1B). The length and width of the galls (1–6 mm and 1–5 mm, respectively) varied with the maturity of the infection (Figs. 1 and 2). Infected *H. stipulacea* fragments were significantly smaller ($p < 0.05$) than uninfected fragments across all biometrics measured (i.e. leaf length, leaf width, internode length, root length) (Fig S1). Mean leaf lengths for healthy and infected *H. stipulacea* fragments were 41.1 mm (95 % CI: 40.1–42.1, $n = 157$) and 22.5 mm (95 % CI: 21.6–23.4, $n = 106$) respectively. Mean leaf widths were 6.25 mm (95 % CI: 6.18–6.32, $n = 157$) and 4.35 mm (95 % CI: 4.25–4.35, $n = 106$) respectively. Mean internode lengths were 9.86 mm (95 % CI: 9.20–10.56, $n = 56$) and 5.56 mm (95 % CI: 5.05–6.07, $n = 43$) respectively, and mean root lengths were 68.3 mm (95 % CI: 63.7–72.8, $n = 36$) and 42.6 mm (95 % CI: 38.9–46.5, $n = 35$) respectively.

Gall coloration and swelling of the leaf petioles of the collected fragments (Fig. 2) were comparable to observations of phytomyxid infected *H. stipulacea* by Vohnik et al. (2017) in the Mediterranean. Based on our observations, we suggest that the infection in *H. stipulacea* fragments on St. Eustatius is caused by a phytomyxid parasite. The morphological changes observed in infected fragments from our study

were not present in the control group of uninfected fragments. Although stunted growth in parts (e.g. internodes, roots, and leaves) of certain seagrass species (e.g. *Zostera noltii* and *Zostera capricorni*) caused by the its specific phytomyxid parasite infection have been observed (Den Hartog, 1989), morphological changes have previously not been described for phytomyxid-infected *H. stipulacea*. It is therefore uncertain from our study to what extent such morphological changes are common in *H. stipulacea*, or which phytomyxid species is causing the infection. Furthermore, phytomyxid-infected *H. stipulacea* was observed during 2018 both in Lac Bay, Bonaire (pers. comm. B. van Tussenbroek) (Fig S2) and Fort Bay, Saba (Maitz, pers. obs.), which are located approximately 810 km south-west and 30 km north-east of St Eustatius, respectively. This suggests that the phytomyxid infecting *H. stipulacea* may have a widespread distribution in the Caribbean. Further examination of the phytomyxid to reveal its phylogenetic and taxonomic affiliation is necessary, as well as understanding the potential ecological effects on its host within its expanding distributional range in the Caribbean.

CRediT authorship contribution statement

Anna Maitz: Conceptualization, Methodology, Investigation, Data curation, Writing - original draft. **Kimani Kitson-Walters:** Investigation, Data curation, Writing - review & editing. **Erik Maitz Boman:** Visualization, Formal analysis, Writing - review & editing.

Declaration of Competing Interest

The authors Anna Maitz, Kimani Kitson-Walters and Erik Maitz Boman declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the

online version, at doi:<https://doi.org/10.1016/j.aquabot.2020.103321>.

References

- Braselton, J.P., 1995. Current status of the plasmodiophorids. *Crit. Rev. Microbiol.* 21, 263–275.
- Braselton, J.P., Short, F.T., 1985. Karyotypic analysis of *Plasmodiophora diplantherae*. *Mycologia* 77, 940–945. <https://doi.org/10.2307/3793306>.
- Den Hartog, C., 1970. The Sea-grasses of the World. North-Holland, Amsterdam.
- Den Hartog, C., 1989. Distribution of *Plasmodiophora bicaudata*, a parasitic fungus on small *Zostera* species. *Diseases of Aquatic Organisms*.
- Ferdinandson, C., Winge, O., 1914. *Ostenfeldia*, a new genus of the Plasmodiophoraceae. *Ann. Bot.* 28, 643–649.
- Fox, J., Weisberg, S., 2018. An R Companion to Applied Regression. Sage publications.
- Karling, J.S., 1968. The Plasmodiophorales, 2nd edn. Hafner Publishing Company, New York.
- Kolátková, V., Čepička, I., Gargiulo, G.M., Vohník, M., 2020. Enigmatic phytomyxid parasite of the alien seagrass *Halophila stipulacea*: new insights into its ecology, phylogeny, and distribution in the Mediterranean Sea. *Microb. Ecol.*
- Levene, H., 1960. Contributions to probability and statistics. *Essays in Honor of Harold Hotelling*, pp. 278–292.
- Maier, I., Parodi, E., Westermeyer, R., Müller, D.G., 2000. *Maulinia ectocarpii* gen. Et sp. nov. (Plasmodiophorea), an intracellular parasite in *Ectocarpus siliculosus* (Ectocarpales, Phaeophyceae) and other filamentous brown algae. *Protist* 151, 225–238.
- Neuhauser, S., Kirchmair, M., Gleason, F.H., 2011. Ecological roles of the parasitic phytomyxids (plasmodiophorids) in marine ecosystems—a review. *Mar. Freshw. Res.* 62, 365–371.
- R Core Team, 2020. R: a Language and Environment for Statistical Computing. URL. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.
- Royston, P., 1995. Remark AS R94: a remark on algorithm AS 181: the W-test for normality. *J. R. Stat. Soc. Ser. C Appl. Stat.* 44, 547–551.
- Ruiz, H., Ballantine, D.L., 2004. Occurrence of the seagrass *Halophila stipulacea* in the tropical west Atlantic. *Bull. Mar. Sci.* 75, 131–135.
- Vohník, M., Borovec, O., Özbek, E.Ö., Aslan, E.Ş.O., 2017. Rare phytomyxid infection on the alien seagrass *Halophila stipulacea* in the southeast Aegean Sea. *Mediterr. Mar. Sci.* 18, 433–442.
- Walker, A.K., Campbell, J., 2009. First records of the seagrass parasite *Plasmodiophora diplantherae* from the northcentral Gulf of Mexico. *Gulf Caribb. Res.* 21, 63–65.
- Wickham, H., 2018. *Francis R. Dplyr: a Grammar of Data Manipulation*. R Package Version 0.4.3. 2015. ed.