

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/378970027>

Social display in the Curaçao Whiptail (*Cnemidophorus murinus* Laurenti)

Article in *Caribbean Herpetology* · March 2024

DOI:10.31611/ch.94

CITATIONS

0

READS

5

1 author:



Philip Senter

Fayetteville State University

110 PUBLICATIONS 1,419 CITATIONS

SEE PROFILE

Social display in the Curaçao Whiptail (*Cnemidophorus murinus Laurenti*)

Philip J. Senter¹ 

¹ Department of Biological and Forensic Sciences, Fayetteville State University, Fayetteville, North Carolina, USA.

Email: psenter@uncfsu.edu

Edited by: R. Graham Reynolds. **Date of publication:** 14 March 2024.

Citation: Senter PJ. 2024. Social display in the Curaçao Whiptail (*Cnemidophorus murinus Laurenti*). *Caribbean Herpetology* 94: 1–4.

DOI: <https://doi.org/10.31611/ch.94>

Three teiid species (Squamata: Teiidae) are endemic to the ABC islands (Aruba, Bonaire, and Curaçao) of the former Netherlands Antilles: the Aruba Whiptail (*Cnemidophorus arubensis*) of Aruba, the Bonaire Whiptail (*C. ruthveni*) of Bonaire and Klein-Bonaire, and the Curaçao Whiptail (*C. murinus*) of Curaçao and Klein-Curaçao. Before now, social displays have been described only for one of these three species: *C. ruthveni* (Baird et al. 2003). All three species engage in an arm-waving display that involves forelimb circumduction (Baird et al. 2003; van Buurt 2005; van Buurt 2011), which conveys social signals to conspecifics in some iguanian and lacertid lizard species (Carpenter et al. 1970; Verbeek 1972; Mitchell 1973). However, this behavior in the ABC island species of *Cnemidophorus* has not been shown to have a social function (Baird et al. 2003). Instead, experimental evidence demonstrates that in *C. ruthveni* it functions as a pursuit deterrent display that is performed in response to approaching humans, whom the lizard may view as potential predators (Cooper et al. 2004), although there are not yet any published reports of *C. ruthveni* performing the display in response to the approach of a non-human predator. My observations and those of others (van Buurt 2011) also confirm that in *C. arubensis* the display is performed in response to approaching humans.

Baird et al. (2003) documented social displays of *C. ruthveni*. These include a challenge display between males, in which a male aggressor dorsally arches his neck and shoulder area while lowering his head, holding his tail on the ground, and laterally compressing his torso. A subsequent display by the challenger and the opponent involves dewlap extension and dorsoventral sigmoidal bobbing of the head and shoulder region while moving slowly forward. In some cases, an aggressor male adds a display in which he turns in a circle with his body flattened to the ground. The courtship display of a male toward a female resembles the challenge display with simultaneous dewlap extension and dorsoventral sigmoidal bobbing of the head and shoulder region while moving toward the female.

The study in which the social displays of *C. ruthveni* were described listed the lizard species as *C. murinus* because, at the time, the Bonaire and Curaçao populations of *Cnemidophorus* were considered to be two subspecies of *C. murinus*. However, the two are now considered distinct species (Ugueto and Harvey 2010). Most that has been published on the natural history of *C. murinus* sensu lato (e.g. Bennett and Gleeson 1979; Bennett and Gorman 1979; Dearing and Schall 1992, 1994; Schall and Dearing 1994; Schall 1996, 2000; Cooper et al. 2003, 2004) pertains to *C. ruthveni*. The natural history of *C. murinus* has been comparatively little studied. Reports have been published on its parasites (Specian and Whittaker 1980), pathology (Hughes and Delis 2014), clutch size (van Buurt 2011), conservation (van Buurt 2006), diet (Senter 2024), and interactions with tourists (van Buurt 2011), but not on its social displays. Van Buurt (2011) illustrated a male *C. murinus* performing a challenge display resembling that of *C. ruthveni*, but no further social displays have been reported in this species before now. Here, I report an observation of social display in *C. murinus*.

This observation occurred between 11.00 and 13.00 h during sunny weather on 29 December 2023, at Playa Kenepa Grandi, Curaçao. The population density of *C. murinus* at Playa Kenepa Grandi is high, and on this day I observed numerous instances of lizards foraging within 50 cm of each other, without conflict and ignoring each other. But I did observe an aggressive encounter between an adult and a juvenile about half the length of

the adult, on a concrete path near the parking lot (12.351111, -69.151667; elevation 8 m asl). The two lizards were moving about on the concrete path, apparently ignoring each other until they were near each other. When the adult was about 50 cm to the juvenile's right, with its anterior end oriented toward the juvenile's right flank, the adult suddenly lunged at the juvenile, without displaying and with a posture typical of locomotion (i.e. without noticeable arching or flattening or other elements typical of display). The juvenile immediately responded by quickly arching its dorsum upward so that its torso was convex dorsally, upon which the adult abruptly stopped the lunge about 10–20 cm from the juvenile. The juvenile then resumed its previous foraging posture, and the two continued to forage without interacting, each moving in a different direction. All of these behaviors, including the lunge, the response, the resumption of normal posture, and the resumption of foraging in different directions, took place in less than 2 s.

It is noteworthy that the display that the juvenile performed is not part of the known repertoire of display behavior in *C. ruthveni*. In the latter, a male challenging another male or courting a female arches its anterior body, but the behavior reported here is thus far the only reported instance of arching of the dorsum as a response to a perceived threat in any species of *Cnemidophorus*, although it has been observed in other teiids (Bostic 1966; Herrel et al. 2009).

Arching of the dorsum is a phylogenetically widespread element of display in lizards, and its known use differs between lizard taxa. It is part of a challenge display between males in numerous gekkotan, lacertid, and iguanian species (Verbeek 1972; Carpenter 1978, 1982; Marcellini 1997) and in the teiid *Aspidoscelis hyperythrus* (Bostic 1966). It is part of a defensive display, directed at humans, in numerous gekkotan species (Marcellini 1997) and in the teiid *Salvator merianae* (Herrel et al. 2009). It has been observed as part of an aggressive, antipredator threat display toward a snake in the phrynosomatid *Sceloporus grammicus* (Pruett et al. 2015). It is part of the male courtship display in several gekkotan species (Marcellini 1997) and is part of a rejection display by females toward males in several iguanian species (Carpenter 1962, 1963, 1967; Ávila and Cunha-Avellar 2006; Vaz e Nunes et al. 2008; Pruett 2014). In *Varanus komodoensis*, it is part of a display by smaller individuals when approaching larger ones, a display that has been termed an appeasement display and which seems to function as the reptilian equivalent of a humble request to exist unmolested nearby (Auffenberg 1978). Displays that appear to serve the same purpose in other lizard species have been termed assertion displays (Carpenter 1967, 1982). They involve arching of the dorsum in the iguanid *Dipsosaurus dorsalis* (Carpenter 1961) and the teiids *Aspidoscelis hyperythrus* and *A. labialis* (Bostic 1966). The behavior in the juvenile *C. murinus* reported here appears to have functioned similarly to the appeasement or assertion displays of other lizard species. Appeasement/assertion displays have not previously been reported for any species of *Cnemidophorus*.

The variety of social displays in *C. ruthveni* (Baird et al. 2003) and the high population densities of its congeners on Aruba and Curaçao suggest that a variety of undocumented social displays might be present in *C. arubensis* and *C. murinus*. *C. murinus* is abundant at Playa Kenepa Grandi and other locations on Curaçao, providing plentiful opportunities to further document other social displays in this species.

References

- Auffenberg W. 1978. Social and feeding behavior in *Varanus komodoensis*. In: Greenberg N, MacLean PB, editors. *Behavior and neurology of lizards: an interdisciplinary colloquium*. Rockville, MD: National Institute of Mental Health. p. 301–331.
- Ávila RW, Cunha-Avellar L. 2006. *Tropidurus etheridgei* (NCN). Courtship. *Herpetological Review* 37: 473.
- Baird TA, Vitt LJ, Baird TD, Cooper WE Jr., Caldwell JP, Pérez-Mellado V. 2003. Social behavior and sexual dimorphism in the Bonaire whiptail, *Cnemidophorus murinus* (Squamata: Teiidae): the role of sexual selection. *Canadian Journal of Zoology* 81: 1791–1790. [Article](#)

- Bennett AF, Gleeson TT. 1979. Metabolic expenditure and the cost of foraging in the lizard *Cnemidophorus murinus*. *Copeia* 1974: 573–577. [Article](#)
- Bennett AF, Gorman GC. 1979. Population density and energetics of lizards on a tropical island. *Oecologia* 42: 339–358. [Article](#)
- Bostic DL. 1966. Threat-display in the lizards *Cnemidophorus hyperythrus* and *Cnemidophorus labialis*. *Herpetologica* 22: 77–79. [Article](#)
- Carpenter CC. 1961. Patterns of social behavior in the desert iguana *Dipsosaurus dorsalis*. *Copeia* 1961: 396–405. [Article](#)
- Carpenter CC. 1962. Patterns of behavior in two Oklahoma lizards. *American Midland Naturalist* 67: 132–151. [Article](#)
- Carpenter CC. 1963. Patterns of behavior in three forms of the fringe-toed lizards (*Uma*-Iguanidae). *Copeia* 1963: 406–412. [Article](#)
- Carpenter CC. 1967. Display patterns of the Mexican iguanid lizards of the genus *Uma*. *Herpetologica* 30: 285–293. [Article](#)
- Carpenter CC. 1978. Ritualistic social behaviors in lizards. In: Greenberg N, MacLean PB, editors. *Behavior and neurology of lizards: an interdisciplinary colloquium*. Rockville, MD: National Institute of Mental Health. p. 253–267.
- Carpenter CC. 1982. The aggressive displays of iguanine lizards. In: Burghardt GM, Rand AS, editors. *Iguanas of the world: their behavior, ecology, and conservation*. Park Ridge, NJ: Noyes. pp. 215–231.
- Carpenter CC, Badham JA, Kimble B. 1970. Behavior patterns of three species of *Amphibolurus* (Agamidae). *Copeia* 1970: 497–505. [Article](#)
- Cooper WE Jr, Pérez-Mellado V, Baird T, Baird TA, Caldwell JP, Vitt LJ. 2003. Effects of risk, cost, and their interactions on optimal escape by nonrefuging Bonaire whiptail lizards, *Cnemidophorus murinus*. *Behavioral Ecology* 14: 288–293. [Article](#)
- Cooper WE Jr, Pérez-Mellado V, Baird TA, Caldwell JP, Vitt LJ. 2004. Pursuit deterrent signalling in the Bonaire whiptail lizard *Cnemidophorus murinus*. *Behaviour* 141: 297–311. [Article](#)
- Dearing MD, Schall JJ. 1992. Testing models of optimal diet assembly by the generalist herbivorous lizard *Cnemidophorus murinus*. *Ecology* 73: 845–858. [Article](#)
- Dearing MD, Schall JJ. 1994. Atypical reproduction and sexual dimorphism of the tropical Bonaire island whiptail lizard *Cnemidophorus murinus*. *Copeia* 1994: 760–766. [Article](#)
- Herrel A, Andrade DV, Eduardo de Carvalho J, Brito A, Abe A, Navas C. 2009. Aggressive behavior and performance in the tegu lizard *Tupinambis merianae*. *Physiological and Biochemical Zoology* 82: 680–685. [Article](#)
- Hughes DF, Delis PR. 2014. *Cnemidophorus murinus murinus* (Laurent's Whiptail). *Myiasis. Herpetological Review* 45: 126–127.
- Marcellini D. 1977. Acoustic and visual display behavior of gekkonid lizards. *American Zoologist* 17: 251–260. [Article](#)
- Mitchell FJ. 1973. Studies on the ecology of the agamid lizard *Amphibolurus maculosus* (Mitchell). *Transactions of the Royal Society of South Australia* 97: 47–76.
- Pruett JA. 2014. *Sceloporus parvus* (Blue-bellied Lizard). Female aggression. *Herpetological Review* 45: 507.
- Pruett JA, Ossip-Klein AG, Vital-Garcia C, Zúñiga-Vega GG, Hews DK. 2015. *Sceloporus grammicus* (Mesquite Lizard). Pursuit-deterrent Signaling. *Herpetological Review* 46: 437–438.
- Schall JJ. 1996. *Cnemidophorus murinus ruthveni* (Bonaire Island Whiptail Lizard). Seeds in diet. *Herpetological Review* 27: 142–143.
- Schall JJ. 2000. Learning in free-ranging populations of the whiptail lizard *Cnemidophorus murinus*. *Herpetologica* 56: 38–45.
- Schall JJ, Dearing MD. 1994. Body temperature of the herbivorous Bonaire Island whiptail lizard (*Cnemidophorus murinus*). *Journal of Herpetology* 28: 526–528. [Article](#)
- Senter PJ. 2024. Diet and foraging in the Curaçao Whiptail, *Cnemidophorus murinus* (Laurenti, 1768). *Caribbean*

- Herpetology* 93: 1–3. [Article](#)
- Specian RD, Whittaker FH. 1980. A survey of the helminth fauna of *Cnemidophorus murinus* from the island of Curacao. *Proceedings of the Helminthological Society of Washington* 47: 275–276.
- Ugueto GN, Harvey MB. 2010. Southern Caribbean *Cnemidophorus* (Squamata: Teiidae): Description of new species and taxonomic status of *C. murinus ruthveni* Burt. *Herpetological Monographs* 24: 111–148. [Article](#)
- van Buurt G. 2005. *Field guide to the amphibians and reptiles of Aruba, Curaçao and Bonaire*. Frankfurt am Main: Chimaira.
- van Buurt G. 2006. Conservation of amphibians and reptiles in Aruba, Curaçao and Bonaire. *Applied Herpetology* 3: 307–321. [Article](#)
- van Buurt G. 2011. The teiid lizards of Aruba, Curaçao, Bonaire (Dutch Caribbean), and the Península de Paraguaná (Venezuela). *Reptiles & Amphibians* 18: 93–104. [Article](#)
- Vaz e Nunes J, Elisei T, Maria de Sousa B. 2008. *Tropidurus itambere* (Rock Lizard). Social interactions. *Herpetological Review* 39: 351–352.
- Verbeek B. 1972. Ethologische Untersuchungen an einigen europäischen Eidechsen. *Bonner Zoologische Beiträge* 23: 122–151. [Article](#)